

Mathematical Symbol Table

Greek			Hebrew		Boldface	Sans Serif	'Blackboard'	Script	Gothic
Name	small	CAPITAL	Name		a	A	À	À	à
Alpha	α	Α	Aleph	א	b	B	Β	Β	ב
Beta	β	Β	Beth	ב	c	C	Ⓒ	Ⓒ	כ
Gamma	γ	Γ	Gimmel	ג	d	D	Ⓓ	Ⓓ	ດ
Delta	δ	Δ	Daleth	ד	e	E	Ⓔ	Ⓔ	Ⓔ
Epsilon	ϵ or ε	Ε			f	F	Ⓕ	Ⓕ	Ⓕ
Zeta	ζ	Ζ			g	G	Ⓖ	Ⓖ	Ⓖ
Eta	η	Η			h	H	Ⓗ	Ⓗ	Ⓗ
Theta	θ or ϑ	Θ			i	I	Ⓘ	Ⓘ	Ⓘ
Iota	ι	Ι			j	J	Ⓙ	Ⓙ	Ⓙ
Kappa	κ	Κ			k	K	Ⓚ	Ⓚ	Ⓚ
Lambda	λ	Λ			l	L	Ⓛ	Ⓛ	Ⓛ
Mu	μ	Μ		Nabla	m	M	Ⓜ	Ⓜ	Ⓜ
Nu	ν	Ν			n	N	Ⓝ	Ⓝ	Ⓝ
Xi	ξ	Ξ			p	P	Ⓟ	Ⓟ	Ⓟ
Omicron	ο	Ο			q	Q	Ⓠ	Ⓠ	Ⓠ
Pi	π or ϖ	Π			r	R	Ⓡ	Ⓡ	Ⓡ
Rho	ρ or ϱ	Ρ			s	S	Ⓢ	Ⓢ	Ⓢ
Sigma	σ or ς	Σ			t	T	Ⓣ	Ⓣ	Ⓣ
Tau	τ	Τ			u	U	Ⓤ	Ⓤ	Ⓤ
Upsilon	υ	Υ			v	V	Ⓤ	Ⓤ	Ⓤ
Phi	ϕ or φ	Φ			w	W	Ⓤ	Ⓤ	Ⓤ
Chi	χ	Χ			x	X	Ⓤ	Ⓤ	Ⓤ
Psi	ψ	Ψ			y	Y	Ⓤ	Ⓤ	Ⓤ
Omega	ω	Ω			z	Z	Ⓤ	Ⓤ	Ⓤ

Logic		Functions							
$\forall x$	'for all ...'	$f : \mathbf{X} \rightarrow \mathbf{Y}$	' f is a function from \mathbf{X} to \mathbf{Y} '						
$\exists x$	'there exists an x such that...'	$f : \mathbf{X} \ni x \mapsto y \in \mathbf{Y}$	' f is a function from \mathbf{X} to \mathbf{Y} ' mapping element x to element y '						
$\exists! x$	'there exists a unique x such that...'	$f : \mathbf{X} \hookrightarrow \mathbf{Y}$	$\mathbf{X} \subset \mathbf{Y}$, and f is the identity map, taking $x \in \mathbf{X}$ to $x \in \mathbf{Y}$						
$\nexists x$	'there does not exist any x ...'	$f : \mathbf{X} \rightarrowtail \mathbf{Y}$	f is an injective function from \mathbf{X} to \mathbf{Y}						
$A \implies B$	'if A , then B ', or, ' A implies B '	$f : \mathbf{X} \twoheadrightarrow \mathbf{Y}$	f is a surjective function from \mathbf{X} to \mathbf{Y}						
$A \Leftarrow B$	'if B , then A ', or, ' B implies A '	Id	The identity map: $\text{Id}(x) = x$ for all x .						
$A \iff B$	' A if and only if B ', or, ' A is equivalent to B '	$\mathbb{1}$	The constant unity: $\mathbb{1}(x) = 1$ for all x .						
TFAE	'The Following Are Equivalent...'	$f^{-1}\{y\}$	\{ $x \in \mathbf{X} ; f(x) = y\}$; the fibre over y or preimage of y (where $f : \mathbf{X} \rightarrow \mathbf{Y}$)						
\square	Q.E.D. —End of Proof.								
↯ or \otimes	Contradiction.								

Set Theory									
$\mathcal{A} \subset \mathcal{B}$	\mathcal{A} is a subset of \mathcal{B} ie. if $a \in \mathcal{A}$, then $a \in \mathcal{B}$ also.					$\mathcal{A} \subseteq \mathcal{B}$	\mathcal{A} is a subset of \mathcal{B} , and possibly $\mathcal{A} = \mathcal{B}$.		
$\mathcal{A} \sqcup \mathcal{B}$	The disjoint union : $\mathcal{A} \sqcup \mathcal{B} = \mathcal{A} \cup \mathcal{B}$, with the assertion that $\mathcal{A} \cap \mathcal{B} = \emptyset$.					$\mathcal{A} \times \mathcal{B}$	The Cartesian product of \mathcal{A} and \mathcal{B} : $\mathcal{A} \times \mathcal{B} = \{(a, b) ; a \in \mathcal{A} \& b \in \mathcal{B}\}$		
$\bigcup_{n=1}^{\infty} \mathcal{A}_n$	$\mathcal{A}_1 \cup \mathcal{A}_2 \cup \mathcal{A}_3 \cup \dots$					$\bigcap_{n=1}^{\infty} \mathcal{A}_n$	$\mathcal{A}_1 \cap \mathcal{A}_2 \cap \mathcal{A}_3 \cap \dots$		
$\bigsqcup_{n=1}^{\infty} \mathcal{A}_n$	$\mathcal{A}_1 \sqcup \mathcal{A}_2 \sqcup \mathcal{A}_3 \sqcup \dots$					$\prod_{n=1}^{\infty} \mathcal{A}_n$	$\mathcal{A}_1 \times \mathcal{A}_2 \times \mathcal{A}_3 \times \dots$		
$\mathcal{A} \setminus \mathcal{B}$	The difference of \mathcal{A} from \mathcal{B} : $\mathcal{A} \setminus \mathcal{B} = \{a \in \mathcal{A} ; a \notin \mathcal{B}\}$					$\mathcal{A} \triangle \mathcal{B}$	The symmetric difference : $\mathcal{A} \triangle \mathcal{B} = (\mathcal{A} \setminus \mathcal{B}) \sqcup (\mathcal{B} \setminus \mathcal{A})$		